

# Please Keep Your Child Home if...



I have a rash or head lice

I'm vomiting



I have diarrhea



I have a fever and another symptom



I don't feel good enough to play

<b>Vomiting</b>	2 or more times in the past 24 hours.
<b>Rash</b>	<ul style="list-style-type: none"> <li>• With fever or behavior change, until a physician has determined the illness is not contagious.</li> <li>• Chickenpox (varicella) until all lesions have dried or crusted.</li> <li>• Impetigo, until 24 hours after treatment has been started.</li> <li>• Mouth sores with drooling.</li> <li>• Head lice or ringworm until after the first treatment.</li> </ul>
<b>Diarrhea</b>	Diapered children whose stool is not contained in the diaper and toilet-trained children with diarrhea causing "accidents" require exclusion from care. In addition, children with diarrhea should be excluded if the stool frequency exceeds 2 or more stools above normal in a day for that child.
<b>Fever</b>	Temperature above 100° F axillary [armpit] or 101° F orally <b>and</b> behavior change or other illness symptoms (e.g., sore throat, rash, vomiting, diarrhea).
<b>Not Feeling Well</b>	Any child who requires more care than can be given by staff in a group setting or a child who is unable to participate in regular daily activities.

For more information on illness exclusion, check with the director or your facility's Health Policy.



**PUBLIC HEALTH** Every One. Every Day. Every Where.